

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/194604>

Please be advised that this information was generated on 2019-06-02 and may be subject to change.

Treatment of Extracranial Carotid Artery Aneurysms: What is Known and What is Unknown?

G.J. de Borst^{*}, C.J.H. van Laarhoven, V.E.C. Pourier

Department of Vascular Surgery, University Medical Centre Utrecht, University Utrecht, Utrecht, The Netherlands

Park and Kim report on a patient with bilateral extracranial carotid artery aneurysms (ECAA) presenting with mild lococervical discomfort but no cerebrovascular sequelae.¹ The ECAA diameter was 2.1 cm on one side and 3 cm on the contralateral side but lengthy over 5 cm. The focus of both the case report and the related commentary was on the surgical intervention performed, but the most important question is if this patient needed intervention at all.²

Park and Kim report that surgical treatment of ECAAs is required in most cases because of the high risk of fatal complications related to embolisation, rupture, and local compression. The authors refer to literature published nearly two decades ago.

This statement is incorrect for two reasons. Firstly, the risk of any ECAA related symptom (including local cervical compression) is probably much lower than previously considered. Prognosis depends on the aetiological background, with post-cervical dissection aneurysms showing a relatively benign course.^{2,3} Secondly, of all symptoms, most are related to local compression, followed by thrombo-embolisation. Rupture of a true ECAA, however, has as far as we are aware, never been reported in the literature.³

Park and Kim state that up to two thirds of cases are related to atherosclerosis; however, probably the largest proportion of ECAAs is related to cervical dissection, either spontaneous or post-traumatic.^{2–4} Pathological examination can make a clear distinction between atherosclerotic ECAA wall versus post-dissection ECAA.⁴ This observation has led to a recently started research program studying vessel wall characteristics to non-invasively predict the risk of ECAA growth over the years.

In the meantime, based on the scarce data available in the literature, an online registry has begun to collect data on patients with ECAA (www.carotidaneurysmregistry.com). The aim of this ongoing international registry is to study the natural course of ECAA and to collect details on interventions for ECAA in a prospective fashion.

Once both the patient and the multidisciplinary team are convinced that intervention is indicated several well described surgical techniques are available. More recently, hybrid interventions using bare metal stents for distally located ECAA were defined,⁵ whereas flow dividing stents may be an option for ECAA at the base of the skull.⁶

Overall, the current literature on treatment outcomes in ECAA consists mainly of case reports and small case series with incomplete data and lack of long-term follow-up.⁷ There is a lack of natural follow-up data and there is no clear treatment algorithm. The online registry to collect data on patients with ECAA is designed to provide clinical guidance on this unusual pathology.

REFERENCES

- 1 Park KM, Kim DI. Bilateral extracranial carotid artery aneurysms treated by staged surgical repair. *EJVES Short Rep* 2017;**36**:13–5.
- 2 Pourier VEC, de Borst GJ. Which carotid artery aneurysms need to be treated (and how)? *J Cardiovasc Surg (Torino)* 2016;**57**(2): 152–7.
- 3 Welleweerd JC, den Ruijter HM, Nelissen BG, Bots ML, Kappelle LJ, Rinkel GJ, et al. Management of ECAA. *Eur J Vasc Endovasc Surg* 2015;**50**(2):141–7.
- 4 Welleweerd JC, Nelissen BG, Koole D, de Vries JP, Moll FL, Pasterkamp G, et al. Histological analysis of extracranial carotid artery aneurysms. *PLoS One* 2015;**10**(1):e0117915.
- 5 Welleweerd JC, de Borst GJ, de Groot D, van Herwaarden JA, Lo RT, Moll FL, et al. Bare metal stents for treatment of extracranial internal carotid artery aneurysms: long-term results. *J Endovasc Ther* 2015;**22**(1):130–4.
- 6 Welleweerd JC, Moll FL, de Borst GJ. Technical options for the treatment of extracranial carotid artery aneurysms. *Expert Rev Cardiovasc Ther* 2012;**10**(7):925–31.
- 7 Welleweerd JC, de Borst GJ. Extracranial carotid artery aneurysm: optimal treatment approach. *Eur J Vasc Endovasc Surg* 2015;**49**(3):235–6.

DOI of original article: <https://doi.org/10.1016/j.ejvssr.2016.11.002>

^{*} Corresponding author. Department of Vascular Surgery G04.129, University Medical Centre Utrecht, PO Box 85500, 3508 GA Utrecht, The Netherlands.

E-mail address: g.j.deBorst-2@umcutrecht.nl (G.J. de Borst).

2405-6553/© 2017 The Author(s). Published by Elsevier Ltd on behalf of European Society for Vascular Surgery. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

<https://doi.org/10.1016/j.ejvssr.2017.11.002>